



Docosahexaenoic acid in Arctic charr (*Salvelinus alpinus*): The importance of dietary supply and physiological response during the entire growth period



David S. Murray^{a,*}, Hannes Hager^a, Douglas R. Tocher^b, Martin J. Kainz^a

^a WasserCluster – Biologische Station Lunz, 3293 Lunz am See, Austria

^b Institute of Aquaculture, School of Natural Sciences, University of Stirling, Stirling, Scotland FK9 4LA, UK

ARTICLE INFO

Article history:

Received 23 July 2014

Received in revised form 2 November 2014

Accepted 7 November 2014

Available online 15 November 2014

Keywords:

Retention

Nutrition

DHA

Aquaculture

ABSTRACT

The aim of this 14-month feeding study was to investigate the effects of dietary docosahexaenoic acid (DHA) on tissue fatty acid composition, DHA retention, and DHA content per biomass accrual in muscle tissues of Arctic charr (*Salvelinus alpinus*). A control feed, formulated with a relatively high DHA inclusion level (F1), was compared with feeds containing gradually reduced amounts of DHA (Feeds F2, F3, and F4). Arctic charr were randomly distributed among 12 tanks and fed one of the feeds in triplicate. The DHA content within muscle tissues of fish fed diets F1 and F2 was generally higher compared to fish fed diets F3 and F4. However, there was an interaction between dietary DHA treatment and season, which resulted in fish muscle tissues having similar DHA contents irrespective of dietary supply during specific sampling periods. Although diets F3 and F4 contained ~4-fold less DHA compared to diets F1 and F2, the retention of DHA in dorsal and ventral muscle tissue was up to 5-fold higher relative to the diet content in fish fed diets F3 and F4. However, the difference among treatments was dependent on the month sampled. In addition, younger fish retained DHA more efficiently compared to older fish. DHA ($\mu\text{g DHA/g/day}$) accrual in muscle tissue was independent of somatic growth, and there was no difference among treatments. The results suggested that dietary DHA may be essential throughout the life cycle of Arctic charr and that the DHA content of muscle tissues was influenced by diet and metabolic/physiological factors, such as specific DHA retention during the entire growth cycle. Finally, this long-term feeding study in Arctic charr indicated a non-linear function in DHA retention in dorsal and ventral muscle tissues throughout the life cycle, which varied in its relationship to dietary DHA.

© 2014 Elsevier Inc. All rights reserved.

1. Introduction

Likely all fish require dietary omega-3 long-chain polyunsaturated fatty acids (n-3 LC-PUFA) as nutrients that function as structural components for cell membranes and facilitate somatic growth and development (Glencross, 2009; Glencross et al., 2014). However, the fatty acid (FA) composition of fish tissues appears to vary widely among species (Glencross, 2009). Studies regarding the importance of specific FA in farmed and wild fish populations suggested that FA compositions of fish tissue lipids usually reflected those of dietary lipids, meaning that tissue FA profiles can be modified by altering the types of dietary fats and oils consumed by fish (Bell et al., 2001; Jobling, 2003, 2004; Sargent et al., 1995). This forms the basis for the use of FA as dietary biomarkers in food web studies (Kirsch et al., 1998) and assumes that the fish do not considerably alter dietary FA.

While it is often assumed that tissue FA reflects dietary FA compositions, recent studies showed that fish metabolism and physiology are also important in influencing the final tissue FA composition in farmed and wild fish. For instance, during FA metabolism many fish, including salmonids such as Atlantic salmon (*Salmo salar*), brown trout (*Salmo trutta*), and Arctic charr (*Salvelinus alpinus*), can convert, albeit rather inefficiently, α -linolenic acid (LNA; 18:3n-3) to the n-3 LC-PUFA eicosapentaenoic acid (EPA; 20:5n-3) and docosahexaenoic acid (DHA; 22:6n-3) (Tocher, 2003; Murray et al., 2014). Furthermore, when analysing FA retention efficiencies, there are distinct differences in the utilisation of different FA (Glencross et al., 2014). For example, previous studies have shown that DHA is preferentially retained in fish tissues when dietary sources are limited (Glencross et al., 2003, 2014; Glencross and Rutherford, 2011; Murray et al., 2014).

The fact that fish can preferentially retain and/or endogenously synthesise DHA may highlight how important this particular LC-PUFA is for somatic development, as all FA can also be β -oxidised to produce metabolic energy (Bell et al., 2002; Tocher, 2003; Sargent et al., 1999). Previous studies have shown that fish fed DHA deficient diets have

* Corresponding author. Tel.: +43 7486 20060 (Office), +43 6802 202422 (Mobile); fax: +43 7486 2006020.

E-mail addresses: david.murray@wcl.ac.at (D.S. Murray), hannes.hager@wcl.ac.at (H. Hager), d.r.tocher@stir.ac.uk (D.R. Tocher), martin.kainz@donau-uni.ac.at (M.J. Kainz).

significantly lower growth rates compared to fish fed diets relatively rich in DHA (Murray et al., 2014).

The majority of studies examining tissue FA compositions focused on the influences of dietary FA on muscle tissue composition and somatic growth (Glencross et al., 2014; Czesny et al., 1999). Furthermore, many studies were performed over relatively short periods of time, typically between 8 and 16 weeks (Turchini et al., 2009) and on different stages of development (Glencross et al., 2003, 2014; Sheridan et al., 1985). Although salmonids are a successfully farmed fish family, DHA requirements are not well defined and information regarding DHA composition in tissues over an entire life cycle is essential in understanding those requirements. However, few studies have thus far investigated how fish accumulate and retain DHA in tissues over an entire life cycle in an effort to elucidate the presumably continuous somatic requirement of this n-3 LC-PUFA.

In the current study, we examined the effect of four dietary treatments containing decreasing dietary DHA on tissue FA compositions, DHA retention ratios, and DHA content per biomass accrual in Arctic charr muscle tissue over an entire production cycle of 14 months. We tested the hypothesis that the mass ratios of DHA (i.e., mg DHA per g muscle tissue) accrued would not differ among fish fed diets containing high and low mass ratios of DHA throughout the life cycle. Our underlying assumption was that fish fed on DHA deficient diets would specifically retain and accrue DHA in muscle tissue and that DHA retention was independent of somatic growth.

2. Materials and methods

2.1. Fish, husbandry, and experimental diets

Third generation farmed Arctic charr (15–20 g body weight) of the same strain (hatchery-reared in Lunz am See, Austria) were held in aquarium facilities at the WasserCluster Research Centre from August 2012 until October 2013. The experiment was conducted in a flow-through system containing 12 rectangular tanks (1000 L each) with a continuous supply of gravel-filtered spring water (ca. 25 L min⁻¹). Waste water was drained using a sink hole covered by a 5 mm mesh screen. Fish were subjected to natural photoperiod (latitude, 47.8604°N), delivered by artificial fluorescent lighting and adjusted weekly. A total of 1200 juvenile Arctic charr were randomly distributed as 100 fish of mixed sexes per tank. Dissolved oxygen, pH, and water temperature were recorded daily. Throughout the long-term feeding trial, Arctic charr was exposed to natural (ambient) water temperature (3.7 °C to 12.3 °C; mean = 7.9 °C), dissolved oxygen (7.3 to 11.4 mg L⁻¹; mean = 9.2 mg L⁻¹) and approximately neutral pH values (6.7 to 7.7; mean = 7.4).

Four isocaloric fish feeds were formulated (Garant™, Austria) to provide sufficient lipid and protein to meet somatic requirements of salmonids (NRC, 2011) (Table 1). Fish in triplicate tanks were fed one

of the four different diets that contained gradually less DHA (Table 2). Diets were dispensed daily into the tank by a clockwork belt feeder (Dryden Aqua Ltd., Edinburgh, UK) over a 12 h feeding period. The daily feed ration exceeded the recommended feeding rate for salmonids for the prevailing water temperature. Uneaten feed was collected to accurately determine feed intake per tank.

2.2. Proximate analysis

The gross nutrient composition of the four experimental diets was determined as described below (Table 3). Moisture was determined by drying to constant weight in an oven at 110 °C for 24 h (Bell et al., 2003). Sample weight was recorded before drying and after removal from the oven. This process was repeated at 1 h intervals until weight change was <5 mg. Total protein content in experimental diets was determined by a modified Bradford assay (Murray et al., 2013) and total lipids by solvent extraction and gravimetric determination (Heissenberger et al., 2010). Ash content was determined by placing pre-weighed diets in a muffle furnace at 550 °C for 8 h or until white ash was obtained (Bell et al., 2003) that was subsequently weighed.

2.3. Sampling procedure

Twelve fish were selected at random, 3 replicates per treatment, and weighed during November, February, April, June, August, and October. Fish were killed by a blow to the head and samples of muscle dissected and stored in plastic vials (8 mL). Muscle samples were obtained by cutting a fillet from the fish and separating the two sections using the lateral line as a border between the dorsal and ventral tissue. Care was taken to prevent any skin or bone from being included in the sample. All tissue samples were stored at -80 °C overnight and freeze dried before analysis.

2.4. Lipid extraction and fatty acid analysis

Total lipids from homogenised and freeze-dried dorsal and ventral muscle samples (25–35 mg) were analysed as by Heissenberger et al. (2010). In brief, samples were sonicated and vortexed in chloroform/methanol (2:1 by volume). Organic layers were removed and transferred into solvent-rinsed vials. For the gravimetric determination of total lipid contents (i.e., mg lipids g dry weight⁻¹), triplicate subsamples (100 µL) of the extracts were evaporated and weighed. Fatty acids were derivatised to obtain fatty acid methyl esters (FAME) using toluene and sulfuric acid-methanol solution (1% v/v, 16 h at 50 °C). In contrast to Heissenberger et al. (2010), hexane without butylated hydroxytoluene (BHT) was used for each washing step after methylation to avoid BHT-related peak interference in chromatograms. FAME

Table 1
Formulations (g/100 g) of experimental feeds F1–F4.

Ingredients	F1	F2	F3	F4
Fish meal, anchovy, super prime, 67% CP	35.0	22.5	22.5	10.0
Pumpkin kernel cake, 59% CP, 11% C. lipids	–	12.5	12.5	25.0
Sunflower protein concentrate, 46% CP	16.8	13.8	13.8	11.0
Haemoglobin powder	7.5	7.5	7.5	7.5
Rapeseed cake, 32.5% CP, 9% CL	5.0	5.0	5.0	5.0
Wheat gluten 80% CP	–	3.34	3.34	6.27
Wheat, feed quality	10.5	9.7	9.7	8.5
Wheat feed flour	6.0	6.0	6.0	6.0
Fish oil (salmon oil)	18.1	17.8	3.0	3.0
Rapeseed oil	–	–	14.8	14.5
Monocalciumphosphate	–	0.6	0.6	1.45
Lysine-HCL	–	0.16	0.16	0.68
Premix	0.8	0.8	0.8	0.8
Diamol (marker)	0.3	0.3	0.3	0.3

Table 2

Polyunsaturated fatty acid (PUFA) contents (mean ± SD; g FA/kg dry weight of diet) of experimental feeds.

Fatty acids	F1	F2	F3	F4
18:3(n-3)	7.4 ± 1.4	7.3 ± 2.7	12.0 ± 1.9	12.0 ± 2.1
18:4(n-3)	0.0 ± 0.0	0.0 ± 0.0	0.0 ± 0.0	0.0 ± 0.0
20:3(n-3)	0.5 ± 0.0	0.5 ± 0.0	0.1 ± 0.0	0.1 ± 0.0
20:4(n-3)	0.0 ± 0.0	0.0 ± 0.0	0.0 ± 0.0	0.0 ± 0.0
20:5(n-3)	8.5 ± 1.8	7.0 ± 1.9	2.8 ± 0.2	2.1 ± 0.7
22:5(n-3)	1.7 ± 0.4	1.6 ± 0.4	0.5 ± 0.0	0.5 ± 0.1
22:6(n-3)	8.4 ± 1.5	7.0 ± 2.7	2.5 ± 0.8	2.0 ± 0.7
Total n-3 PUFA	26.5 ± 5.1	23.4 ± 7.7	17.8 ± 2.9	16.7 ± 3.7
18:2(n-6)	23.8 ± 11.0	25.1 ± 2.9	36.4 ± 1.0	38.9 ± 3.2
18:3(n-6)	0.2 ± 0.1	0.2 ± 0.0	0.0 ± 0.0	0.0 ± 0.0
20:2(n-6)	3.1 ± 1.2	2.4 ± 0.7	1.0 ± 0.0	1.1 ± 0.4
20:3(n-6)	0.3 ± 0.1	0.3 ± 0.0	0.1 ± 0.0	0.1 ± 0.1
20:4(n-6)	0.7 ± 0.1	0.5 ± 0.2	0.2 ± 0.0	0.2 ± 0.0
22:4(n-6)	0.1 ± 0.0	0.1 ± 0.0	0.0 ± 0.0	0.0 ± 0.0
Total n-6 PUFA	30.0 ± 12.9	30.4 ± 4.2	38.5 ± 1.1	41.0 ± 4.1

Table 3
Proximate composition of experimental diets (g/100 g of diet).

	F1	F2	F3	F4
Protein	43.2 ± 1.0	43.7 ± 2.4	44.6 ± 2.1	44.0 ± 4.0
Lipid	25.1 ± 2.3	24.5 ± 1.4	24.4 ± 1.1	23.8 ± 3.4
Ash	10.2 ± 1.3	8.4 ± 0.0	8.0 ± 0.1	8.5 ± 0.9
Moisture	7.2 ± 0.3	5.8 ± 0.3	8.1 ± 0.3	8.8 ± 1.3

were identified by comparison with known standards (Supelco37 FAME Mix) using a gas chromatograph (Thermo Scientific TRACE GC Ultra™) equipped with a flame ionisation detector (FID) and a Supelco™ SP-2560 column (100 m, 25 mm i.d., 0.2 µm film thickness). Quantification of FA was performed by comparison with a known concentration of the internal standard using Excalibur 1.4™ (Thermo Electron Corporation).

2.5. Data analysis

Significant differences among dietary treatments were determined by one-way ANOVA and interactions between treatment and season were determined using general linear models (GLM). Differences between means were determined by Tukey's HSD test. Relationships between retention ratios and somatic growth was analysed using linear regressions. Data identified as nonhomogeneous, using variance test, were subjected to log transformation before applying the statistical tests. The Minitab®16 statistical software package was used for data analysis and the significance level was set at $P = 0.05$. Feed conversion ratios (FCR) were determined as the quotient of feed consumed and increase in fish biomass, feed intake (g)/weight gain (g). Fatty acid retention ratios were determined as the quotient of FA in fish muscle tissues and FA in the respective diet (mg FA per g of dry weight/diet); i.e., $[DHA]_{fish}/[DHA]_{diet}$. We define retention as the ability of fish to regulate and control ingested FA in their tissues (Kainz et al., 2004). Fatty acid biomass accrual was determined as the quotient of measured FA over time (mg FA per g of dry weight/time) i.e., $[DHA]_{fish}/time(days)$. Furthermore, we define DHA biomass accrual as the DHA content in muscle tissue per weight of fish, which was normalised to account for differences in somatic growth (i.e., DHA per unit biomass accrual).

3. Results

3.1. Diet composition, fish growth, and feed utilisation

Dietary lipid content was similar among treatments. There was a 1.6-fold decrease in total n-3 PUFA contents between diets F1 and F4, specifically a 4.0 and 4.2-fold decrease in the proportions DHA and EPA, respectively (Table 2). Alternatively, total n-6 PUFA contents increased 1.4-fold and LNA by 1.6-fold between diets F1-F4 (Table 2).

Dietary treatment had a significant effect on final weight ($F_{[3-11]} = 26.70$; $R^2 = 0.875$; $P < 0.001$) (Table 4). Fish fed dietary treatment F1 (350.0 ± 22.8 g/fish) and F2 (291.9 ± 1.4 g/fish) had significantly higher final weights compared to fish fed F4 (236.3 ± 16.9 g/fish) diets. Weight gain of fish among treatments was also significantly different ($F_{[3-11]} = 26.41$; $R^2 = 0.874$; $P < 0.001$) (Table 4). Fish fed dietary treatments F1 and F2 gained significantly more weight

(338.8 ± 22.5 g/fish and 281.6 ± 12.2 g/fish, respectively) compared to fish fed diet F4 (226.0 ± 17.1 g/fish). Significant differences in FCR were also observed during the experiment ($F_{[3-92]} = 26.41$; $R^2 = 0.501$; $P < 0.001$). Fish fed diet F1 (0.91 ± 0.04) had significantly lower FCR compared to fish fed diet F2 (1.02 ± 0.05 , F3 (1.06 ± 0.09) and F4 (1.13 ± 0.11) (Table 4). Regression analysis showed no linear relationship between fish weight and dietary or tissue DHA contents.

3.2. Effect of seasonality on tissue DHA content in fish fed different diets

There were no significant differences in DHA contents of dorsal or ventral muscle among fish fed the different dietary treatments at the end of the feeding experiment (Fig. 1). Seasonality interacted significantly with diet treatments for DHA in dorsal muscle ($F_{[15-71]} = 2.12$; $R^2 = 0.728$; $P = 0.025$) (Fig. 1). In February, fish fed diet F1 had significantly higher DHA (61.2 ± 11.0 mg DHA/g total lipids; $F_{[3-11]} = 6.52$; $R^2 = 0.710$; $P = 0.015$) compared to fish fed dietary treatment F4 (25.2 ± 10.2 mg DHA/g total lipids). In April, dorsal muscle DHA in fish fed diets F1 (126.1 ± 30.3 mg DHA/g total lipids) and F2 (148.7 ± 13.8 mg DHA/g total lipids) were significantly higher ($F_{[3-11]} = 9.34$; $R^2 = 0.778$; $P = 0.005$) compared to fish fed diet F4 (56.0 ± 17.2 mg DHA/g total lipids). In August, fish fed diets F1 (137.8 ± 6.3 mg DHA/g total lipids) and F2 (125.4 ± 32.3 mg DHA/g total lipids) also had significantly higher DHA in dorsal muscle ($F_{[3-11]} = 33.09$; $R^2 = 0.925$; $P < 0.001$) compared to fish fed diets F3 (22.8 ± 17.7 mg DHA/g total lipids) and F4 (20.8 ± 8.4 mg DHA/g total lipids). There were, however, no significant effects of dietary treatment on DHA contents of dorsal muscle in fish sampled during November, June or October (Fig. 1).

Similarly, seasonality showed a significant interaction with dietary treatment in ventral muscle DHA ($F_{[15-71]} = 2.03$; $R^2 = 0.749$; $P = 0.033$) (Fig. 2). In April, ventral muscle DHA contents were significantly higher in fish fed diets F1, F2, and F3 (118.6 ± 14.1 , 120.4 ± 39.5 , and 95.7 ± 19.0 mg DHA/g total lipid, respectively; $F_{[3-11]} = 10.79$; $R^2 = 0.802$; $P = 0.003$) compared to fish fed diet F4 (38.5 ± 15.8 mg DHA/g total lipids). There were no significant differences in DHA ventral muscle contents among fish sampled during November, February, June, August, or October (Fig. 2).

Dorsal and ventral muscle DHA contents in fish fed diets F3 and F4 were generally, but not significantly lower compared to fish fed diets F1 and F2. The seasonal patterns of DHA content in dorsal muscle were similar between November and June independent of dietary treatment (Fig. 1); fish fed diets F1, F2, and F3 showed a 3-fold increase in DHA in dorsal muscle, while DHA content doubled in fish fed diet F4. Between June and October, the pattern of DHA between fish fed high (F1 and F2) and low (F3 and F4) DHA diets diverged; by August, the DHA content in fish fed diets F1 and was 6- to 7-fold higher than in fish fed diets F3 and F4, while it converged again by October at the end of the trial. In general, DHA in ventral muscle tissue followed a similar pattern independent of dietary treatment (Fig. 2). Between February and April, there was a large increase in DHA with fish fed diets F1 and F2 diets showing levels increased 4- and 2-fold, respectively, and fish fed diets F3 and F4 showing increases of 2- and 1.5-fold, respectively (Fig. 2). From April to October, DHA levels increased 3-fold in fish fed diets F1 and F2, and 2-fold in fish fed diets F3 and F4 (Fig. 2).

Table 4
Growth and feed utilisation (mean ± SD) of Arctic charr fed different diets over the 401 day experimental period.

	F1	F2	F3	F4
Initial weight (g fish ⁻¹)	11.1 ± 0.3	10.3 ± 0.5	10.4 ± 0.7	10.3 ± 0.9
Final weight (g fish ⁻¹)	350.0 ± 22.8 ^a	291.9 ± 1.4 ^b	270.3 ± 8.0 ^{bc}	236.3 ± 16.9 ^c
Weight gain (g fish ⁻¹)	338.8 ± 22.5 ^a	281.6 ± 12.2 ^b	259.9 ± 8.4 ^{bc}	226.0 ± 17.1 ^c
Total feed deposited in tanks (kg/tank)	20.1 ± 1.1 ^a	16.9 ± 0.6 ^b	14.7 ± 1.5 ^b	16.0 ± 1.1 ^b
Feed conversion ratio	0.91 ± 0.05 ^a	1.02 ± 0.05 ^b	1.06 ± 0.09 ^b	1.13 ± 0.11 ^c




Fig. 1. Dorsal muscle DHA (mean \pm SD) contents (mg/g total lipids) of Arctic charr fed diets containing different DHA contents. Significant differences are indicated above relevant months.

3.3. DHA in muscle tissues per biomass accrual

There was no significant difference in DHA contents in dorsal (Fig. 3) or ventral (Fig. 4) muscle tissues per biomass accrual among the different dietary treatments. Both tissues follow similar patterns independent of dietary DHA supply with highest DHA accrual ($\mu\text{g DHA/g/day}$) at the beginning of the trial in juvenile Arctic charr (November) with a sharp decrease (~ 4 -fold in dorsal and ventral muscle tissues) until February, followed by consistent increase until April, but with generally lower DHA content in both ventral and dorsal muscle tissues per biomass accrual until the end of the trial.

3.4. Effect of seasonality on DHA retention ratios

Retention ratios of DHA in dorsal ($F_{[3-11]} = 28.10$; $R^2 = 0.913$; $P < 0.001$) and ventral ($F_{[3-11]} = 38.98$; $R^2 = 0.936$; $P < 0.001$) muscle tissue differed significantly at the end of the feeding experiment (October). In

dorsal muscle, fish fed diets F3 (3.3 ± 0.8) and F4 (3.3 ± 0.2) had significantly higher retention ratios compared to fish fed diets F1 (1.1 ± 0.3) and F2 (1.3 ± 0.2) (Fig. 5). In ventral muscle tissues, fish fed diets F3 (4.5 ± 0.9) and F4 (3.3 ± 0.4) also had significantly higher retention ratios compared to fish fed diets F1 (1.0 ± 0.1) and F2 (1.3 ± 0.4) (Fig. 6). There was no significant interaction between season and dietary treatment in DHA retention ratios of dorsal or ventral muscle tissues. However, differences in DHA retention ratios between dietary treatments all varied on a monthly basis (Figs. 5 and 6).

In dorsal muscle from fish sampled during November, DHA retention in fish fed diet F4 (4.6 ± 1.5) was significantly higher ($F_{[3-11]} = 5.14$; $R^2 = 0.530$; $P = 0.029$) compared to fish fed diet F1 (3.8 ± 0.7) (Fig. 5). Retention of DHA in April was also significantly higher ($F_{[3-11]} = 12.65$; $R^2 = 0.826$; $P = 0.002$) in fish fed diets F3 (4.3 ± 0.6) and F4 (5.5 ± 3.0) compared to fish fed diets F1 (1.2 ± 0.1) and F2 (1.5 ± 0.1) (Fig. 5). There were no significant differences in DHA retention ratios in dorsal muscle tissue during February, June, or August (Fig. 5).




Fig. 2. Ventral muscle DHA (mean \pm SD) contents (mg/g total lipids) of Arctic charr fed diets containing different DHA contents. Significant differences are indicated above relevant months.




Fig. 3. DHA per unit biomass accrual in dorsal muscle tissue ($\mu\text{g DHA/g/day}$).

Retention of DHA in ventral muscle of fish fed diets F3 (3.6 ± 0.4) and F4 (4.6 ± 1.5) was significantly higher ($F_{[3-11]} = 8.23$; $R^2 = 0.755$; $P = 0.008$) than in fish fed diet F1 (1.3 ± 0.3) during November (Fig. 6). In February, fish fed diets F3 (2.3 ± 0.1) and F4 (2.5 ± 0.4) also retained significantly higher ($F_{[3-11]} = 6.12$; $R^2 = 0.697$; $P = 0.018$) levels of DHA in ventral muscle than fish fed diet F1 (0.6 ± 0.5) (Fig. 6). In June, fish fed diets F3 (2.0 ± 0.1) and F4 (2.5 ± 1.2) had significantly higher ($F_{[3-11]} = 8.93$; $R^2 = 0.770$; $P = 0.006$) DHA retention ratios compared to fish fed diets F1 (0.9 ± 0.4) and F2 (0.8 ± 0.0) (Fig. 6). Ventral muscle DHA retention ratios in fish fed diets F3 (1.9 ± 1.0) and F4 (2.6 ± 0.5) were significantly higher ($F_{[3-11]} = 7.39$; $R^2 = 0.735$; $P = 0.011$) than fish fed diet F2 (0.4 ± 0.2) in August (Fig. 6).

3.5. Relationships between DHA retention ratios in muscle tissue and weight of Arctic charr

DHA retention in dorsal ($R^2 = 0.22$; $P < 0.001$) and ventral ($R^2 = 0.16$; $P = 0.001$) muscle tissues decreased significantly from juvenile to adult fish (Figs. 7 and 8, respectively). The linear relationship between DHA retention in muscle tissue and fish weight depended on dietary treatments. In dorsal muscle, there was a significant negative relationship between fish weight and DHA retention ratios for fish fed diets F1 ($R^2 = 0.31$, $P = 0.017$), F3 ($R^2 = 0.30$, $P = 0.019$) and F4 ($R^2 = 0.22$, $P = 0.048$), but not fish fed diet F2 (Fig. 7). In ventral muscle, there was also a significant negative relationship between fish weight and DHA retention ratio for fish fed diets F2 ($R^2 = 0.28$, $P = 0.022$) and F3 ($R^2 = 0.27$, $P = 0.027$), but not fish fed diets F1 and F4 (Fig. 8).

4. Discussion

This study demonstrated that both dietary DHA supply and fish physiology during the entire growth cycle affect DHA retention




Fig. 4. DHA per unit biomass accrual in ventral muscle tissue ($\mu\text{g DHA/g/day}$).

efficiency and, consequently, DHA contents in dorsal and ventral muscle tissues of Arctic charr. One possible explanation underpinning DHA retention in fish and the higher DHA content compared to all other FA is that DHA is not easily β -oxidised as the $\Delta 4$ double bond requires peroxisomal oxidation (Sargent et al., 2002; Tocher, 2003; Stillwell and Wassall, 2003). However, this biochemical mechanism based on enzyme specificities and activities may not be the only one involved, and the present study has suggested that other processes may also interact to affect DHA retention. Final weights and weight gain in fish fed low amounts of dietary DHA (F3–F4) were lower compared to fish supplied with higher amounts of DHA (F1–F2). Furthermore, FCR was also highest in fish supplied with the lowest amount of DHA (F4). In addition, fish fed lower dietary DHA had correspondingly generally lower DHA contents in muscle tissues, but higher DHA retention ratios compared to fish supplied with higher dietary DHA. The similar DHA content per unit biomass accrual was largely independent of dietary DHA supply, indicating that increased DHA retention in Arctic charr compensates, at least partly, for dietary deficits. However, this effect appeared more pronounced in younger fish as evidenced by the decreased DHA retention ratios in muscle tissues in fish of increasing size and thus age. Furthermore, the present study also showed that, independent of dietary DHA supply, DHA in muscle tissues fluctuated during the life cycle of Arctic charr, which suggests that DHA metabolism in Arctic charr may change with extrinsic factors, such as water temperature (Farkas et al., 1980), as well as intrinsic factors, such as gonadal development (Hiratsuka et al., 2004). This long-term study has thus provided better resolution of DHA retention in tissues over time compared to shorter feeding studies that can miss seasonal fluctuations.

The present study demonstrated that diets containing lower amounts of DHA resulted in lower final weights and reduced weight gain, particularly with the lowest inclusion levels (diet F4) compared to fish fed F1 and F2 diets. In addition, FCR was lower in fish fed the F1 diet compared to all other dietary treatments. These results are in contrast to previous studies which showed no significant impact of lower dietary DHA on growth rates and/or final fish weights (Gomes et al., 1995; Guillou et al., 1995; Kaushik et al., 1995; Petterson et al., 2009) and no significant differences in FCR of farmed fish (Azevedo et al., 2004; Bendiksen et al., 2003; Karalazos et al., 2011). Many studies examining the effect of fatty acid content on final weights and weight gain were performed over relatively short time scales (Turchini et al., 2009). For example, studies reporting no significant difference in growth rates between fish consuming high and low amounts of DHA were performed between 12 and 21 weeks (Bell et al., 2001, 2003; Tocher et al., 2000, 2001; Torstensen et al., 2000).

However, the current study showed no significant relationship between DHA and fish weight. A previous study found that, after 50 wks of feeding, Atlantic salmon fed low amounts of DHA (2.9 and 3.4 g/100 g of total fatty acids) had significantly higher weights compared to fish fed higher amounts of DHA (10.5 g/100 g of total fatty acids) (Bell et al., 2003). Therefore, the lower growth rates in fish fed diets F2–F4 in the current study may have been a consequence of the replacement of fish meal with pumpkin kernel cake. Pumpkins contain high contents of neutral and acid detergent fibre (Saura-Calixto et al., 1983) that affect digestive functions by increasing intestinal flow rates (Liener 1969, Krogdahl, 1989; Nyina-Wamwiza et al., 2010), which may reduce the retention of dietary nutrients (Krogdahl, 1989). This suggested that a 2-fold increase in pumpkin kernel cake in the present study may have affected nutrient absorption, resulting in the lower growth rates of fish fed dietary treatments F2, F3 and F4.

Dietary DHA supply did not fully predict the DHA content in dorsal or ventral muscle tissues of Arctic charr. There were significant interactions between dietary treatment and season in both muscle tissues. In general, DHA content was higher in fish fed F1 and F2 diets, but these results depended on the month sampled. Previous studies have shown that dietary FA requirements for fish differ depending on life history




Fig. 5. DHA retention ratios in dorsal muscle of Arctic charr during the feeding period.

stage (Izquierdo, 1996; Izquierdo et al., 2001); thus, we may expect that tissue FA compositions in fish vary similarly reflecting the different biological requirements associated with each stage of their life cycle. Sheridan et al. (1985) found significant differences in FA in muscle tissues between different life history stages of steelhead trout (*Oncorhynchus mykiss*) fed the same diet. The DHA content of smolt muscle tissue was significantly higher compared to parr muscle tissue. This suggested that fish may require different amounts of DHA at different stages of development to support crucial, albeit not clearly defined, biological functions.

Despite a four-fold decrease in DHA mass ratios between F1/F2 and F3/F4 diets, there were no significant differences in DHA per unit biomass accrual among treatments in both dorsal and ventral muscle tissues. It is known that DHA, with its long carbon chain length and high number of double bonds, provides cell membranes with high flexibility, which improves cellular functions (Arts and Kohler, 2009; Eldho et al., 2003; Wiegand, 1996). Indeed, Arctic charr provided with the lowest dietary DHA generally showed the highest retention ratios in both dorsal and ventral muscle tissues, indicating that in the absence of or reduced dietary DHA, Arctic charr may preferentially retain and/or

synthesise DHA. Turchini and Francis (2009) stated that high deposition of DHA in tissues, in fish fed low dietary DHA, may be due to endogenous synthesis of DHA. Murray et al. (2014) found that there was a trend towards higher DHA synthesis in Arctic charr fed diets containing low DHA. Such endogenous synthesis of DHA may be required to support cell and other physiological functions when dietary DHA supply is low.

The DHA mass ratios retained generally mirrors the DHA content in Arctic charr muscle tissue. For example, the DHA content in muscle tissue of fish fed F3 and F4 diets was high (i.e., April), and retention ratios of DHA were high. Previous studies also report that Atlantic salmon (Glencross et al., 2014) and European sea bass (*Dicentrarchus labrax*) (Mourete and Bell, 2006) have substantially higher DHA retention ratios in tissues when fed diets containing low or no DHA. These studies investigated this process over a much shorter time scale (9–20 weeks) compared to the current study (60 weeks), which found that the degree to which the retention ratios differed among treatments changed depending on the month sampled and thus on the specific stage of somatic fish development. For example, although DHA retention ratios in fish fed diets F3 and F4 were generally higher, there was no significant difference in dorsal muscle DHA retention ratios among treatments in




Fig. 6. DHA retention ratios in ventral muscle of Arctic charr during the feeding period.




Fig. 7. Relationships between DHA retention ratios in dorsal muscle and fish weight, grouped by dietary treatment with corresponding P-value from regression analysis.

February, June, and August. These results indicated that, although fish require high DHA throughout their life cycle, their ability to retain DHA within muscle tissue fluctuated on a month-by-month basis.

Juvenile Arctic charr retained more DHA in dorsal and ventral muscle tissue compared to adult charr. However, this relationship was dependent on dietary treatment. The somatic requirement for DHA is higher in juvenile and sub-adult fish when a large portion of DHA, supporting rapid growth, is directed towards the formation of cell membranes vital for normal development and functioning of tissues (Izquierdo, 1996; Mourente et al., 1991; Stillwell and Wassall, 2003; Tocher et al., 1985). DHA in muscle mostly originates from dietary supplies and, in part, from endogenous synthesis from shorter chain precursors in liver (Tocher, 2003). Therefore, the present results suggested that juvenile Arctic charr are more efficient at retaining dietary DHA and/or the amount of DHA synthesised *de novo* is higher compared to adult Arctic charr. Tocher et al. (2003) found that endogenous synthesis of DHA was higher in juvenile Atlantic salmon (parr) compared to older Atlantic salmon (smolt). During later stages of development, fish allocate energy to gonadal maturation (Bell et al., 1997; Izquierdo et al., 2001; Mourente and Odriozola, 1990; Wiegand, 1996), and thus the retention of DHA in somatic tissues may be reduced in sexually mature adults.

In conclusion, the present study suggests that dietary inclusion of DHA affects both the content and the retention of DHA in dorsal and ventral muscle tissues. However, over the course of an entire life cycle, the significance of these differences varied suggesting that other factors, such as age and stage of sexual development, may also influence the content and retention of DHA in somatic tissues. Although there was a




Fig. 8. Relationship between DHA retention ratios in ventral muscle and fish weight, grouped by dietary treatment with corresponding P-value from regression analysis.

trend towards lower DHA in muscle tissue of fish provided with 4-fold lower dietary DHA, these Arctic charr were able to retain as much as 5-fold more DHA in muscle tissues compared to fish fed the highest dietary DHA inclusion. Furthermore, despite differences in dietary DHA levels, DHA accrual was similar among dietary treatments independent of somatic growth, but varied over time. Therefore, the results suggested that dietary DHA was not the sole predictor of DHA in muscle tissues of Arctic charr and that the DHA content of diets could be assessed based on the ability of the species to specifically retain and/or endogenously synthesise DHA, the age of the fish, and the stage of sexual development.

Acknowledgements

This work received financial support by the Austrian Ministry of Life (project no. 100837; BMLFUW-LE.1.3.2/0051-II/1/2012) and GARANT Tiernahrung Austria. We are grateful for technical assistance and support from Eduard Schneeberger and veterinary supervision by Heinz Heistingner. We also thank Katharina Drucker, Katharina Hader, Katharina Winter, and Zahra Changizi for their laboratory assistance.

References

- Arts, M.T., Kohler, C.C., 2009. Health and Condition in Fish: The Influence of Lipids on Membrane Competency and Immune Response, Lipids in Aquatic Ecosystems. Springer, pp. 237–256.
- Azevedo, P.A., Leeson, S., Cho, C.Y., Bureau, D.P., 2004. Growth, nitrogen and energy utilization of juveniles from four salmonid species: diet, species and size effects. *Aquaculture* 234, 393–414.
- Bell, J.G., Farndale, B.M., Bruce, M.P., Navas, J.M., Carillo, M., 1997. Effects of broodstock dietary lipid on fatty acid compositions of eggs from sea bass (*Dicentrarchus labrax*). *Aquaculture* 149, 107–119.
- Bell, J.G., McEvoy, J., Tocher, D.R., McGhee, F., Campbell, P.J., Sargent, J.R., 2001. Replacement of fish oil with rapeseed oil in diets of Atlantic salmon (*Salmo salar*) affects tissue lipid compositions and hepatocyte fatty acid metabolism. *J. Nutr.* 131, 1535–1543.
- Bell, J.G., Henderson, R.J., Tocher, D.R., McGhee, F., Dick, J.R., Porter, A., Smullen, R.P., Sargent, J.R., 2002. Substituting fish oil with crude palm oil in the diet of Atlantic salmon (*Salmo salar*) affects muscle fatty acid composition and hepatic fatty acid metabolism. *J. Nutr.* 132, 222–230.
- Bell, J.G., Tocher, D.R., Henderson, R.J., Dick, J.R., Crampton, V.O., 2003. Altered fatty acid compositions in Atlantic salmon (*Salmo salar*) fed diets containing linseed and rapeseed oils can be partially restored by a subsequent fish oil finishing diet. *J. Nutr.* 133, 2793–2801.
- Bendiksen, E.A., Berg, O.K., Jobling, M., Arnesen, A.M., Måsøvalb, K., 2003. Digestibility, growth and nutrient utilisation of Atlantic salmon parr (*Salmo salar* L.) in relation to temperature, feed fat content and oil source. *Aquaculture* 224, 283–299.
- Czesny, S., Kolkowski, S., Dabrowski, K., Culver, D., 1999. Growth, survival, and quality of juvenile walleye *Stizostedion vitreum* as influenced by n-3 HUFA enriched *Artemia nauplii*. *Aquaculture* 178, 103–115.
- Eldho, N.V., Feller, S.E., Tristram-Nagle, S., Polozov, I.V., Gawrisch, K., 2003. Polyunsaturated docosahexaenoic vs docosapentaenoic acid differences in lipid matrix properties from the loss of one double bond. *J. Am. Chem. Soc.* 125, 6409–6421.
- Farkas, T., Csengeri, I., Majoros, F., Olah, J., 1980. Metabolism of fatty acids in fish. III. Combined effect of environmental temperature and diet in formation and deposition of fatty acids in the carp, *Cyprinus carpio* Linnaeus 1758. *Aquaculture* 20, 29–40.
- Glencross, B.D., 2009. Exploring the nutritional demand for essential fatty acids by aquaculture species. *Rev. Aquac.* 1, 71–124.
- Glencross, B.D., Rutherford, N.R., 2011. The docosahexaenoic acid (DHA) requirements of juvenile barramundi (*Lates calcarifer*). *Aquac. Nutr.* 17, 536–548.
- Glencross, B.D., Hawkins, W.E., Curnow, J.G., 2003. Restoration of the fatty acid composition of red seabream (*Pagrus auratus*) using a fish oil finishing diet after grow-out on plant oil based diets. *Aquac. Nutr.* 9, 409–418.
- Glencross, B., Tocher, D., Matthew, C., Gordon Bell, J., 2014. Interactions between dietary docosahexaenoic acid and other long-chain polyunsaturated fatty acids on performance and fatty acid retention in post-smolt Atlantic salmon (*Salmo salar*). *Fish Physiol. Biochem.* 40, 1213–1227.
- Gomes, E.d.F., Rema, P., Kaushik, S.J., 1995. Replacement of fish meal by plant proteins in the diet of rainbow trout (*Oncorhynchus mykiss*): digestibility and growth performance. *Aquaculture* 130, 177–186.
- Guillou, A., Soucy, P., Khalil, M., Adambounou, L., 1995. Effects of dietary vegetable and marine lipid on growth, muscle fatty acid composition and organoleptic quality of flesh of brook charr (*Salvelinus fontinalis*). *Aquaculture* 136, 351–362.
- Heissenberger, M., Watzke, J., Kainz, M., 2010. Effect of nutrition on fatty acid profiles of riverine, lacustrine, and aquaculture-raised salmonids of pre-alpine habitats. *Hydrobiologia* 650, 243–254.
- Hiratsuka, S., Kitagawa, T., Matsue, Y., Hashidume, M., Wada, S., 2004. Lipid class and fatty acid composition of phospholipids from the gonads of skipjack tuna. *Fish. Sci.* 70, 903–909.

- Izquierdo, M.S., 1996. Essential fatty acid requirements of cultured marine fish larvae. *Aquac. Nutr.* 2, 183–191.
- Izquierdo, M.S., Fernandez-Palacios, H., Tacon, A.G.J., 2001. Effect of broodstock nutrition on reproductive performance of fish. *Aquaculture* 197, 25–42.
- Jobling, M., 2003. Do changes in Atlantic salmon, *Salmo salar* L., fillet fatty acids following a dietary switch represent wash-out or dilution? Test of a dilution model and its application. *Aquac. Res.* 34, 1215–1221.
- Jobling, M., 2004. 'Finishing' feeds for carnivorous fish and the fatty acid dilution model. *Aquac. Res.* 35, 706–709.
- Kainz, M., Arts, M.T., Mazumder, A., 2004. Essential fatty acids in the planktonic food web and their ecological role for higher trophic levels. *Limnol. Oceanogr.* 49, 1784–1793.
- Karalazos, V., Bendiksen, E.A., Dick, J.R., Tocher, D.R., Bell, J.G., 2011. Influence of the dietary protein:lipid ratio and fish oil substitution on fatty acid composition and metabolism of Atlantic salmon (*Salmo salar*) reared at high water temperatures. *Br. J. Nutr.* 105, 1012–1025.
- Kaushik, S.J., Cravedi, J.P., Lalles, J.P., Sumpter, J., Fauconneau, B., Laroche, M., 1995. Partial or total replacement of fish meal by soybean protein on growth, protein utilization, potential estrogenic or antigenic effects, cholesterolemia and flesh quality in rainbow trout, *Oncorhynchus mykiss*. *Aquaculture* 133, 257–274.
- Kirsch, P.E., Iverson, S.J., Bowen, W.D., Kerr, S.R., Ackman, R.G., 1998. Dietary effects on the fatty acid signature of whole Atlantic cod (*Gadus morhua*). *Can. J. Fish. Aquat. Sci.* 55, 1378–1386.
- Krogdahl, Å., 1989. Alternative protein sources from plants contain antinutrients affecting digestion in salmonids, The current status of fish nutrition in aqua-culture. Proceedings of the third international symposium on feeding and nutrition in fish. August, pp. 253–261.
- Liener, I., 1969. Protease inhibitors. In: Liener, I. (Ed.), Toxic constituents of plant foodstuffs. Elsevier, pp. 7–68.
- Mourente, G., Bell, J.G., 2006. Partial replacement of dietary fish oil with blends of vegetable oils (rapeseed, linseed and palm oils) in diets for European sea bass (*Dicentrarchus labrax* L.) over a long term growth study: Effects on muscle and liver fatty acid composition and effectiveness of a fish oil finishing diet. *Comp. Biochem. Physiol. B Biochem. Mol. Biol.* 145, 389–399.
- Mourente, G., Odriozola, J.M., 1990. Effect of broodstock diets on lipid classes and their fatty acid composition in eggs of gilthead sea bream (*Sparus aurata* L.). *Fish Physiol. Biochem.* 8, 93–101.
- Mourente, G., Tocher, D.R., Sargent, J.R., 1991. Specific accumulation of docosahexaenoic acid (22:6n-3) in brain lipids during development of juvenile turbot *Scophthalmus maximus* L. *Lipids* 26, 871–877.
- Murray, D.S., Bain, M.M., Adams, C.E., 2013. Adhesion mechanisms in European whitefish *Coregonus lavaretus* eggs: is this a survival mechanism for high-energy spawning grounds? *J. Fish Biol.* 83, 1221–1233.
- Murray, D.S., Hager, H., Tocher, D.R., Kainz, M., 2014. Effect of partial dietary fish meal and fish oil replacement pumpkin kernel cake and rapeseed oil on fatty acid composition and metabolism in Arctic charr. (*Salvelinus alpinus*). *Aquaculture* 431, 85–91.
- Nyina-Wamwiza, L., Wathélet, B., Richir, J., Rollin, X., Kestemont, P., 2010. Partial or total replacement of fish meal by local agricultural by-products in diets of juvenile African catfish (*Clarias gariepinus*): growth performance, feed efficiency and digestibility. *Aquac. Nutr.* 16, 237–247.
- Petterson, A., Pickova, J., Brännäs, E., 2009. Effects of crude rapeseed oil on lipid composition in Arctic charr *Salvelinus alpinus*. *J. Fish Biol.* 75, 1446–1458.
- Sargent, J., Bell, G., McEvoy, L., Tocher, D.R., Estevez, A., 1999. Recent developments in the essential fatty acid nutrition of fish. *Aquaculture* 177, 191–199.
- Sargent, J.R., Bell, J.G., Bell, M.V., Henderson, R.J., Tocher, D.R., 1995. Requirement criteria for essential fatty acids. *J. Appl. Ichthyol.* 11, 183–19.
- Sargent, J.R., Tocher, D.R., Bell, J.G., 2002. The Lipids. In: Halver, J.E., Hardy, R.W. (Eds.), *Fish Nutrition*, 3rd edition Academic Press, San Diego, pp. 181–257 (Ch.4).
- Saura-Calixto, F., Canellas, J., Garcia-Raso, J., 1983. Determination of hemicellulose, cellulose and lignin contents of dietary fibre and crude fibre of several seed hulls. Data comparison. *Z. Lebensm. Unters. Forsch.* 177, 200–202.
- Sheridan, M.A., Allen, W.V., Kerstetter, T.H., 1985. Changes in the fatty acid composition of steelhead trout, *Salmo gairdneri* Richardson, associated with parr-smolt transformation. *Comp. Biochem. Physiol. B Biochem. Mol. Biol.* 80, 671–676.
- Stillwell, W., Wassall, S.R., 2003. Docosahexaenoic acid: membrane properties of a unique fatty acid. *Chem. Phys. Lipids* 126, 1–27.
- Tocher, D.R., 2003. Metabolism and functions of lipids and fatty acids in teleost fish. *Rev. Fish. Sci.* 11, 107–184.
- Tocher, D.R., Fraser, A.J., Sargent, J.R., Gamble, J.C., 1985. Lipid class composition during embryonic and early larval development in Atlantic herring (*Clupea harengus*, L.). *Lipids* 20, 84–89.
- Tocher, D.R., Bell, J.G., Dick, J.R., Henderson, R.J., McGhee, F., Michell, D., Morris, P.C., 2000. Polyunsaturated fatty acid metabolism in Atlantic salmon (*Salmo salar*) undergoing parr-smolt transformation and the effects of dietary linseed and rapeseed oils. *Fish Physiol. Biochem.* 23, 59–73.
- Tocher, D.R., Agaba, M., Hastings, N., Bell, J.G., Dick, J.R., Teale, A.J., 2001. Nutritional regulation of hepatocyte fatty acid desaturation and polyunsaturated fatty acid composition in zebrafish (*Danio rerio*) and tilapia (*Oreochromis niloticus*). *Fish Physiol. Biochem.* 24, 309–320.
- Tocher, D.R., Bell, J.G., McGhee, F., Dick, J.R., Fonseca-Madrigal, J., 2003. Effects of dietary lipid level and vegetable oil on fatty acid metabolism in Atlantic salmon (*Salmo salar* L.) over the whole production cycle. *Fish Physiol. Biochem.* 29, 193–209.
- Torstensen, B., Lie, O., Froyland, L., 2000. Lipid metabolism and tissue composition in Atlantic salmon (*Salmo salar* L.)- Effects of capelin oil, palm oil, and oleic acid-enriched sunflower oil as dietary lipid sources. *Lipids* 35, 653–664.
- Turchini, G.M., Francis, D.S., 2009. Fatty acid metabolism (desaturation, elongation and β -oxidation) in rainbow trout fed fish oil-or linseed oil-based diets. *Br. J. Nutr.* 102, 69–81.
- Turchini, G.M., Torstensen, B.E., Ng, W.-K., 2009. Fish oil replacement in finfish nutrition. *Rev. Aquac.* 1, 10–57.
- Wiegand, M.D., 1996. Composition, accumulation and utilization of yolk lipids in teleost fish. *Rev. Fish Biol. Fish.* 6, 259–286.